



# Sepsis in Obstetric Care Change Package



## Authors

**Melissa Bauer, DO**, Faculty, IHI

**Kelly McCutcheon Adams, LICSW**, Senior Director, IHI

**Lauren Enright**, Project Manager, IHI

## Acknowledgments

The Sepsis in Obstetric Care Change Package was made possible with the generous funding of the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number UC4MC28042, Alliance for Innovation on Maternal Health (AIM). We thank Deborah Bamel, MPH and Catherine Mather, MA for their contributions as members of the Institute for Healthcare Improvement (IHI) core team. We also thank our AIM colleagues, Emily Greenwood and Hannah Kaufman, for their steadfast partnership in this work, as well as our numerous expert faculty who contributed during the foundational expert meeting, interviews, and/or subsequent draft reviews: April Chavez; Carol Burke, APRN, CNS, RNC-OB, C-EFM; Catherine Albright, MD, MS; Chelsea Lennox, MPH; Kristin DeVries, MA, MPP; Maile Le Beouf; Malavika Prabhu, MD; Megan Snair, MPH; Phillips Heine, MD; Ronald Gibbs, MD; Tiffany Messerall, DNP, APRN, WHNP-BC, RNC-OB; Rachel Solnick, MD, MSc; and Courtney Salley.

---

These materials were developed with support from the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) as part of a cooperative agreement with the American College of Obstetricians and Gynecologists under grant number UC4MC28042, Alliance for Innovation on Maternal Health. The contents do not necessarily represent the official views of, nor an endorsement, by HRSA, HHS, or the U.S. Government. For more information, please visit [HRSA.gov](https://www.hrsa.gov)

©2022 American College of Obstetricians and Gynecologists. Permission is hereby granted for duplication and distribution of this document, in its entirety and without modification, for solely non-commercial activities that are for educational, quality improvement, and patient safety purposes. All other uses require written permission from ACOG.

# Contents

<b>Introduction</b>	<b>4</b>
Why is this important?	4
What is a change package?	4
How to prioritize changes?	5
<b>Change Package</b>	<b>6</b>
A Note on Symbols	6
Readiness	7
Recognition and Prevention	11
Response	14
Reporting and Systems Learning	17
Respectful, Equitable, and Supportive Care	19
<b>Appendix</b>	<b>21</b>



# Introduction

The Alliance for Innovation on Maternal Health (AIM) is a national data-driven maternal safety and quality improvement (QI) initiative. AIM works through state and community-based teams to align national, state, and hospital-level QI efforts to reduce preventable maternal mortality and severe morbidity across the United States.

The AIM Patient Safety Bundles are a core part of this work. To promote the successful implementation of these bundles, AIM partnered with the Institute for Healthcare Improvement (IHI) to create a series of associated change packages. This specific change package is designed to support Perinatal Quality Collaboratives (PQCs) and other state-based initiatives to leverage the AIM Sepsis in Obstetric Care Patient Safety Bundle more effectively.

## Why is this important?

Obstetric sepsis remains a leading cause of maternal mortality in the United States and throughout the world.<sup>1</sup> Maternal deaths due to sepsis have been found to be largely preventable with timely recognition, appropriate treatment, and escalation of care.<sup>2</sup> Sepsis disproportionately affects those from underrepresented minority groups. National rates of obstetric sepsis are 2.4 times higher for Black patients, 1.5 times higher for Asian/Pacific Islander patients, and 1.8 times higher for Native American patients compared with White patients.<sup>3</sup> These differences directly reflect the effects of racism on maternal morbidity and mortality.

This change package is meant to provide teams with the necessary steps for preparation to recognize, treat, and escalate care to improve outcomes through implementation of the AIM Sepsis in Obstetric Care Patient Safety Bundle.<sup>4</sup> The change ideas and culture shifts focusing on equitable and respectful care outlined in this document will serve to improve patient-centered care not only for obstetric patients with sepsis but for all obstetric patients.

## What is a change package?

A change package is a document listing evidence-based or best-practice changes specific to a topic and is usually organized around a framework or model. In this case, the Sepsis in Obstetric Care Change Package is structured around the [Sepsis in Obstetric Care Patient Safety Bundle](#).<sup>4</sup>

Changes packages, including this one, are structured around the following components:

- **Primary Drivers:** Major processes, operating rules, or structures that will contribute to moving toward the aim. In this change package, the primary drivers are based on AIM's Five Rs Framework (Readiness, Recognition & Prevention, Response, Reporting/Systems Learning, and Respectful Care).
- **Change Concepts:** Broad concepts (e.g., "move steps in the process closer together") that are not yet specific enough to be actionable but that will be used to generate specific ideas for change.

- **Change Ideas:** Actionable, specific ideas for changing a process. Change ideas can come from research, best practices, or from other organizations that have recognized a problem and have demonstrated improvement on a specific issue related to that problem.

Taken as a whole, a change package has the potential to seem overwhelming. Based on the priorities of your state and community, we encourage you to start small by testing a couple of ideas connected to the aim you set. Through iterative tests of change (also known as Plan-Do-Study-Act (PDSA) cycles), you will have an opportunity to learn what works and what does not in your efforts to improve your processes. Initially, these cycles are carried out on a small scale (e.g., one patient on one day) to see if they result in improvement. Teams can then expand the tests and gradually incorporate larger and larger samples until they are confident that the changes will result in sustained improvement.

## How to prioritize changes?

No team is expected to test all the listed change ideas. Consider this a menu of options from which you may choose what to tackle first. Each team will review their baseline data, progress to date, organizational priorities, and select an area(s) to prioritize. For example, some may start with one driver. Others may start by tackling one idea across all drivers. Start by choosing an area that you think could lead to an easy win.


You can also leverage the following tools to help you decide where to start:

1. [Pareto chart](#): A type of bar chart in which the various factors that contribute to an overall effect are arranged in order according to the magnitude of their effect. This ordering helps identify the "vital few" — the factors that warrant the most attention.<sup>5</sup>
2. [Priority matrix](#): A tool that can better help you to understand important relationships between two groupings (e.g., steps in a process and departments that conduct that step) and make decisions on where to focus.<sup>6</sup>
3. [Impact-effort matrix](#): A tool that helps identify which ideas seem easiest to achieve (least effort) with the most effects (highest impact). The ideas identified via this tool would be a great place to start.<sup>7</sup>


# Change Package

## A Note on Symbols

### Respectful, Equitable, and Supportive Care

In the latest revision of the AIM Sepsis in Obstetric Care Patient Safety Bundle, a fifth R was added; Respectful, Equitable, and Supportive Care. This R is integrated throughout the change package, and all change ideas that fall under this R are marked with a  symbol.

### Additional Considerations

It is understood that every team utilizing this change package will be at a different point in this work. If your organization is further along in your obstetric sepsis improvement work and has found reliability in some of the change ideas below, we suggest testing the additional considerations in *italics* and marked by the  symbol.

## Readiness

Change Concept	Change Idea	Key Resources and Tools
<p><b>Establish inter- and intradepartmental protocols and policies for the care of patients experiencing obstetric sepsis or suspected sepsis</b></p>	<p>Establish a reliable and efficient system to order, obtain, and promptly administer appropriate antimicrobials</p> <p><i>Have antibiotics in an automated medication dispensing system with a reliable system to monitor expiration date *</i></p> <p><i>Implement "code sepsis" to alert pharmacy to immediately dose, prepare, and deliver antibiotics to the bedside *</i></p>	<p><a href="#">Risk Factors, Etiologies, and Screening Tools for Sepsis in Pregnant Women: A Multicenter Case-Control Study</a><sup>8</sup></p> <p><a href="#">California Maternal Quality Care Collaborative (CMQCC): Improving Diagnosis and Treatment of Maternal Sepsis: A Quality Improvement Toolkit, Part III. Assessment and Treatment of Maternal Sepsis: Antibiotics and Source Control</a><sup>9</sup></p>
	<p>Create an obstetric-focused, multidisciplinary rapid-response team with the ability to consult at the bedside and recognize/respond to rapid deterioration</p>	<p><a href="#">AIM: Sepsis in Obstetrical Care: Element Implementation Details</a><sup>10</sup></p>
	<p>In hospitals without 24/7 in-hospital obstetrician coverage, establish "first responder" protocol for when sepsis is suspected</p>	<p><a href="#">AIM: Sepsis in Obstetrical Care: Element Implementation Details</a><sup>10</sup></p>
	<p>Create an obstetric-specific protocol for suspected or identified sepsis that includes notification of the obstetric rapid-response team, order sets for prompt treatment (antimicrobials, fluid administration, and vasopressor use), and bedside evaluation for escalation of care. Include criteria for when a patient must be seen in person by an MD or certified nurse midwife (CNM).</p> <p><i>Involve patients with lived experience in development of protocol</i> ♦ *</p>	<p><a href="#">CMQCC: Improving Diagnosis and Treatment of Maternal Sepsis: A Quality Improvement Toolkit, Part III. Assessment and Treatment of Maternal Sepsis: Antibiotics and Source Control</a><sup>9</sup></p>

	<p>Establish protocol for when an obstetrician consult is needed for patients who are &lt;20 weeks pregnant or postpartum</p>	<p><a href="#">Sepsis Bundle Project: National Hospital Inpatient Quality Measures</a><sup>11</sup></p> <p><a href="#">Internal Validation of the Sepsis in Obstetrics Score to Identify Risk of Morbidity From Sepsis in Pregnancy</a><sup>12</sup></p>
<p><b>Provide multidisciplinary education on obstetric sepsis to all clinicians and staff that provide care to pregnant and postpartum people, including in non-labor &amp; delivery settings such as emergency departments, intensive care units, and outpatient clinics</b></p>	<p>Require routine team training based on real-life scenarios, regardless of level of experience/education</p> <p>Include sepsis scenarios in regular unit simulations and drills for obstetric emergencies to engage learners in immediate recognition and treatment of sepsis (e.g., can use “mystery shopper” model where mock patient calls front desk complaining of weakness/shortness of breath)</p> <p>Provide didactic education (such as multi-departmental grand rounds) about obstetric sepsis and related policies, procedures, and teamwork skills</p> <p>Conduct education at orientation, whenever changes to processes or procedures occur, and every two years</p> <p>Ensure that ED, outpatient clinics, nurse advice line, front-desk staff, and any other “entry points” are included in education efforts</p> <p>Include education on pregnancy-specific presentation and treatment of sepsis</p>	<p><a href="#">AIM: Urgent Maternal Warning Signs</a><sup>13</sup></p> <p><a href="#">The UK Sepsis Trust: How to Spot Sepsis</a><sup>14</sup></p>

		<p><a href="#">End Sepsis: What is Sepsis?</a><sup>15</sup></p> <p><a href="#">Sepsis Alliance: Pregnancy &amp; Childbirth</a><sup>16</sup></p> <p><a href="#">Top 10 Pearls for the Recognition, Evaluation, and Management of Maternal Sepsis</a><sup>17</sup></p>
<p><b>Utilize evidence-based criteria for sepsis assessment for all pregnant and postpartum patients, in all units, including obstetric-specific criteria, when appropriate</b></p>	<p>Utilize a sepsis screening tool on admission throughout hospitalization to identify patients who may develop sepsis</p> <p>To avoid missing patients:</p> <ul style="list-style-type: none"> <li>• Use pregnancy-adjusted tool &gt;20 weeks pregnancy and &lt;3 days postpartum</li> <li>• Use a non-pregnancy-adjusted tool for early pregnancy (&lt;20 weeks) and &gt;3 days postpartum</li> </ul>	<p><a href="#">Sepsis Bundle Project: National Hospital Inpatient Quality Measures</a><sup>11</sup></p> <p><a href="#">CMQCC: Improving Diagnosis and Treatment of Maternal Sepsis: A Quality Improvement Toolkit</a><sup>9</sup></p> <p><a href="#">American College of Obstetricians and Gynecologists (ACOG) District II: Maternal Safety Bundle for Sepsis in Pregnancy (slides 22 – 23)</a><sup>18</sup></p> <p><a href="#">United Kingdom Obstetric Surveillance System (UKOSS)</a><sup>19</sup></p>
<p><b>Create a culture that utilizes non-hierarchical communication so that all team members, including the patient, feel empowered to speak up about a concern and know that their input is valued by the entire care team</b></p>	<p>Empower nurses to speak up when they have concerns, and respect nurses' concerns when they do speak up</p> <hr/> <p>Empower patients to speak up when they have concerns and thoroughly evaluate their concerns</p>	<p><a href="#">IHI: Three Ways to Create Psychological Safety in Health Care</a><sup>20</sup></p> <p><a href="#">IHI White Paper: A Framework for Safe, Reliable, and Effective Care</a><sup>21</sup></p> <hr/> <p><a href="#">Centers for Disease Control and Prevention (CDC): Hear Her Campaign</a><sup>22</sup></p>

	<p>Empower physicians to feel safe saying, "I don't know, but I will find out" in team communications</p>	<p><a href="#">IHI: Three Ways to Create Psychological Safety in Health Care</a><sup>20</sup></p> <p><a href="#">IHI White Paper: A Framework for Safe, Reliable, and Effective Care</a><sup>21</sup></p>
	<p>Conduct team training in communicating across power dynamics</p>	<p><a href="#">Agency for Healthcare Research and Quality (AHRQ) TeamSTEPPS</a><sup>23</sup></p>
	<p>Take the time to ask patients questions, listen to their concerns, and better understand their specific context <span style="color: orange;">◇</span></p> <ul style="list-style-type: none"> <li>• Suggestion: sit down so you are at eye level with patient prior to asking about concerns</li> </ul>	

## Recognition and Prevention

Change Concept	Change Idea	Key Resources and Tools
<b>Implement evidence-based measures to prevent infection</b>	Implement a comprehensive list of evidence-based practices to reduce risk of surgical-site infection	<a href="#">ACOG Practice Bulletin Number 195: Prevention of Infection After Gynecologic Procedures</a> <sup>24</sup>
	Ensure prevention measures address pelvic and non-pelvic categories in all phases of care	
	Build pregnancy-specific vital sign change alerts into electronic health record (EHR) systems for >20 weeks of gestation and <3 days postpartum. Use non-pregnant criteria <20 weeks of gestation and >3 days postpartum	<a href="#">CMQCC: Improving Diagnosis and Treatment of Maternal Sepsis: A Quality Improvement Toolkit</a> <sup>9</sup> <a href="#">Severe Maternal Sepsis in the UK, 2011–2012: A National Case-Control Study</a> <sup>25</sup> <a href="#">The maternal early warning criteria: a proposal from the national partnership for maternal safety</a> <sup>26</sup> <a href="#">Use of Maternal Early Warning Trigger tool reduces maternal morbidity</a> <sup>27</sup>
	Consider concerns of patients, doulas, and identified support network to be critical information for further evaluation ♦	
	Consider antibiotic treatment in case of isolated fever during labor, unless another source can be identified	<a href="#">Committee Opinion No. 712 Summary: Intrapartum Management of Intraamniotic Infection</a> <sup>28</sup>

	Utilize order sets for suspected obstetric sepsis	<a href="#">CMQCC: Improving Diagnosis and Treatment of Maternal Sepsis: A Quality Improvement Toolkit<sup>9</sup></a>
	<p>Conduct routine screening for asymptomatic bacteriuria (to prevent maternal progression to pyelonephritis), group B streptococci (GBS) colonization, and sexually transmitted infections during prenatal care (to prevent fetal infection)</p> <ul style="list-style-type: none"> <li>Evaluate at the time of admission for unresolved infections and/or need for GBS prophylaxis</li> </ul>	
<b>Consider sepsis on the differential diagnosis of a person with deteriorating status, even in the absence of fever</b>	Ask follow-up questions to determine severity to nonspecific symptoms that could be misattributed to common challenges during pregnancy and postpartum (such as postsurgical pain, fatigue, and anxiety)	<a href="#">Rutgers Robert Wood Johnson Medical School: Stop. Look. Listen!<sup>29</sup></a>
	Reframe to a high-acuity, low-occurrence mindset to increase suspicion of sepsis	
<b>In all care environments, assess and document if a patient presenting is pregnant or has been pregnant within the past year</b>	Incorporate assessment, ideally in EHR system, in all entrance portals of care (such as ED and urgent care), for patients aged 10 – 55	
	Train associates/departments on the “why” behind screening for current or recent pregnancy	
	Ensure gender inclusivity in assessment ♦	
<b>Provide patient education focused on general life-threatening pregnancy and postpartum complications and early warning signs, including sepsis signs and</b>	<p>Standardize discharge education for patient and their identified support network</p> <ul style="list-style-type: none"> <li>Include standardized education in child-birthing classes and prenatal appointments</li> <li>Use teach-back to assess understanding</li> </ul>	<p><a href="#">AIM: Urgent Maternal Warning Signs<sup>13</sup></a></p> <p><a href="#">CDC: Urgent Maternal Warning Signs<sup>30</sup></a></p>

<p><b>symptoms other than fever, and instructions for who to notify with concerns</b></p>		<p><a href="#">The UK Sepsis Trust: How to Spot Sepsis</a><sup>14</sup></p> <p><a href="#">End Sepsis: What is Sepsis?</a><sup>15</sup></p> <p><a href="#">Sepsis Alliance: Pregnancy &amp; Childbirth</a><sup>16</sup></p> <p><a href="#">Association of Women’s Health, Obstetric, and Neonatal Nurses (AWHONN): Save Your Life Poster</a><sup>31</sup> to be used in conjunction with the <a href="#">AWHONN Post-Birth Warning Signs Education Program</a><sup>32</sup></p>
	<p><i>Engage community-based organizations in development of culturally appropriate and language-specific materials</i> ♦ *</p>	
	<p>Ensure patient education materials are aligned with patients’ health literacy, culture, language, and accessibility needs ♦</p>	<p><a href="#">CDC: Urgent Maternal Warning Signs</a><sup>30</sup></p> <p><a href="#">Centers for Disease Control and Prevention (CDC): Hear Her Campaign</a><sup>22</sup></p> <p><a href="#">Sepsis Alliance: Pregnancy &amp; Childbirth</a><sup>16</sup></p>

## Response

Change Concept	Change Idea	Key Resources and Tools
<p><b>Initiate facility-wide standard protocols and policies for assessment, treatment, and escalation of care for people with suspected or confirmed obstetric sepsis</b></p>	<p>Immediately triage patients when they present with sepsis symptoms</p>	<p><a href="#">AIM: Sepsis in Obstetrical Care: Element Implementation Details</a><sup>10</sup></p>
	<p>Reduce time to initiate appropriate antimicrobials</p>	
	<p>Ensure all antibiotics can be started within one hour concurrently rather than in tandem, with closed-loop communication with nursing and physician leadership to start additional IVs as needed</p>	
	<p>Perform source control urgently when indicated using the least invasive means</p>	
	<p>Use ideal body weight to calculate resuscitation fluids (30mL/kg is likely 1-2 liters)</p>	<p><a href="#">University of Washington: Identification and Management of Sepsis in Pregnancy: Obstetric Consensus Statement</a><sup>33</sup></p>
	<p>Start norepinephrine after adequate fluid resuscitation to avoid fluid overload (if needed to maintain MAP &gt;65 mm Hg)</p>	<p><a href="#">Society for Maternal-Fetal Medicine Consult Series #47: Sepsis during pregnancy and the puerperium</a><sup>34</sup></p>
	<p>Call for support and/or consultation from services such as Infectious Disease, ICU, and Maternal-Fetal Medicine</p>	
	<p>Include “equity pause” to look at bias risk within multidisciplinary care planning and to ask, “What are considerations to ensure respectful care without discrimination?” ♦</p> <p><i>Consider race, ethnicity, language, gender identity, obesity, mental health history, unplanned pregnancy and history of pregnancies,</i></p>	

	<p><i>marital status, housing status, education level, etc. as potential areas of bias for providers</i> ♦ *</p>	
	<p>Designate provider to take lead on patient and family communication during a crisis and ensure use of interpreter when needed ♦</p>	
<p><b>Initiate facility-wide standard protocols and policies for post-stabilization management of people with sepsis</b></p>	<p>Identify hospital maternal level of care and determine if the hospital has ICU capabilities for pregnant or postpartum patients. If no capabilities exist, create a pathway for transfer to an appropriate facility.</p>	<p><a href="#">AIM: Sepsis in Obstetrical Care: Element Implementation Details</a><sup>10</sup></p> <p><a href="#">Levels of Maternal Care: Obstetric Care Consensus No. 9</a><sup>35</sup></p> <p><a href="#">ACOG Practice Bulletin No. 106: Intrapartum fetal heart rate monitoring: nomenclature, interpretation, and general management principles</a><sup>36</sup></p>
	<p>Coordinate daily multidisciplinary team communication, and for any major changes in clinical care, and have a designated team member provide updates to the patient and designated support network</p>	
<p><b>Facilitate comprehensive post-sepsis care, including screening and proper referrals for post-sepsis syndrome</b></p>	<p>Provide mental health support post-diagnosis/discharge ♦</p> <ul style="list-style-type: none"> <li>• Perform anxiety, depression, and PTSD screening at regular intervals</li> <li>• Provide HRSA maternal mental health hotline</li> </ul>	<p><a href="#">Sepsis Alliance: Post-Sepsis Syndrome</a><sup>37</sup></p> <p><a href="#">ACOG Committee Opinion Number 757: Screening for Perinatal Depression</a><sup>38</sup></p> <p><a href="#">HRSA: Primary Care PTSD Screen for DSM-F (PC-PTSD-5)</a><sup>39</sup></p> <p><a href="#">Massachusetts General Hospital (MGH) Center for Women’s Mental Health: Screening for Perinatal Anxiety Using PASS – the</a></p>

		<a href="#">Perinatal Anxiety Screening Scale<sup>40</sup></a> <a href="#">HRSA Maternal &amp; Child Health: National Maternal Mental Health Hotline<sup>41</sup></a>
	Provide information about post-sepsis syndrome and include identified support network in education upon discharge ♦	<a href="#">Sepsis Alliance: Post-Sepsis Syndrome<sup>37</sup></a> <a href="#">PostICU Resources<sup>42</sup></a>
	Screen and, if applicable, refer to rehabilitation care (occupational therapy (OT), physical therapy (PT), and/or speech therapy)	<a href="#">Enhancing Recovery from Sepsis: A Review<sup>43</sup></a>
	Coordinate referrals for continuing multidisciplinary care, such as wound care or nutrition, and include social work for support	

## Reporting and Systems Learning

Change Concept	Change Idea	Key Resources and Tools
<p><b>Conduct multidisciplinary reviews for systems improvement of each sepsis case to assess the screening program, the quality of care provided to patients with sepsis, and whether instances of bias may have impacted care</b></p>	<p>Conduct formal after-action review with designated leader and standardized content</p> <p><i>Reflect on the equity in case as part of review</i> ♦ *</p>	
	<p>Review time from alert to ordering antibiotics, time to start of administration, and time for all antibiotics to be infused. Identify barriers and address them.</p>	
	<p>Share findings from reviews with all associated staff and facility stakeholders at regular staff meetings</p>	
	<p><i>Larger hospital systems can consider having trained personnel from the risk department in dialogue with patients during review process, speaking on the patients’ behalf during internal multidisciplinary review</i> *</p>	
	<p>Consider processes of care and patient outcomes in the context of race, ethnicity, and social determinants of health ♦</p>	<p><a href="#">Developing Tools to Report Racism in Maternal Health for the CDC Maternal Mortality Review Information Application (MMRIA): Findings from the MMRIA Racism &amp; Discrimination Working Group</a><sup>44</sup></p>
	<p>Create a systematic process to screen patient chart language for instances of bias (i.e., “refused” vs. “declined,” using race to describe a patient, using adjective “difficult,” etc.) ♦</p>	<p><a href="#">Negative Patient Descriptors: Documenting Racial Bias In The Electronic Health Record</a><sup>45</sup></p>
	<p>Implement huddles, specifically in the following circumstances:</p> <ul style="list-style-type: none"> <li>• Concern for any patient condition</li> </ul>	

<p><b>Establish a culture of multidisciplinary planning, huddles, and post-event debriefs</b></p>	<ul style="list-style-type: none"> <li>• Worsening of patient condition</li> <li>• Patient thinks concerns are not being heard</li> </ul>	
	<p>Establish standardized briefing documentation to capture successes and determine actionable follow-up</p>	
	<p>Archive debriefing documentation for obstetric sepsis and review systematically with unit-specific and QI leadership teams</p>	
	<p>Establish unit-specific and QI leadership teams to review and address quality and safety issues</p>	
	<p>Conduct post-event debriefing with an equity lens for support and learning ◊</p>	
	<p>Create systemic plan to follow up on bad experiences noted in standard surveys</p> <p><i>Consider Patient Reported Experience Measures (PREM) for patients to complete *</i></p>	
	<p>Include patients and their identified support networks in rounds ◊</p>	
	<p>Regularly clarify what goals and values for care are essential to include in a patient’s treatment plan ◊</p>	

## Respectful, Equitable, and Supportive Care

Change Concept	Change Idea	Key Resources and Tools
<p><b>Include each pregnant or postpartum person and their identified support network as respected members of and contributors to the multidisciplinary care team</b></p>	<p>Engage patients and their care team in decision making about their care at every point, from admission through discharge, including during rounds ♦</p>	
<p><b>Engage in open, transparent, and empathetic communication with pregnant and postpartum people and their identified support network about sepsis diagnosis and recommended treatment plans that are aligned with their health literacy, culture, language, and accessibility needs</b></p>	<p>Do not minimize patient complaints or concerns; explore them with follow-up questions and assessment ♦</p>	<p><a href="http://rutgers.edu">Stop. Look. Listen! (rutgers.edu)</a></p>
	<p>Educate clinicians on providing respectful care by engaging in the life-long learning of cultural humility, understanding that individuals cannot learn all aspects of any culture, including their own ♦</p>	
	<p>Provide communication in the patient’s preferred language ♦</p>	
	<p>Provide educational materials for patients in common languages spoken in your community ♦</p>	
	<p>Engage with patients in open, honest, and respectful communication ♦</p>	<p><a href="http://HospitalCareers.com">15 Bedside Manner Techniques to Improve Patient Experience   HospitalCareers.com</a></p>
	<p>Ask patients if they would like to be accompanied by their support network for any exams, procedures, and discussions ♦</p>	
	<p>Ask patients how they would prefer to be addressed, get pronunciation correct, and share with the entire care team ♦</p>	

<p><b>Because maternal mortality and severe maternal morbidity related to sepsis disproportionately affect Black, Indigenous, and Hispanic people because of systemic racism, but not race itself, it is necessary to mitigate this bias by having a high index of suspicion for sepsis</b></p>	<p>Create system where staff can actively seek feedback on their presentation of biases and be open to correction ◇</p>	
	<p>Set aims for closing identified disparities using the SMARTIE format (strategic, measurable, ambitious, realistic, time-bound, inclusive, and equitable) ◇</p>	<p><a href="#">IHI: Science of Improvement: Setting Aims<sup>46</sup></a></p> <p><a href="#">IHI: Science of Improvement: Tips for Setting Aims<sup>47</sup></a></p> <p><a href="#">IHI: Applying an Equity Lens to Performance Improvement<sup>48</sup></a></p>
	<p>Collect and analyze REAL (Race, Ethnicity, and Language) data ◇</p> <p><i>Have staff training on importance of REAL data and respectful collection ◇ *</i></p>	
	<p>Review all process and outcome data disaggregated by REAL to assess for inequities with unit-specific and QI leadership teams ◇</p> <p><i>Engage leaders in messaging about destigmatizing discussion and identification of inequities to move toward action ◇ *</i></p> <p><i>Identify alternative strategies to integrate equity considerations into reporting and systems learning in settings where use of disaggregated data may cause potential patient identifiability or unstable data ◇ *</i></p>	
	<p>Assess quality of REAL data and develop processes for improved data collection ◇</p> <p><i>Identify a champion focused on inequities ◇ *</i></p>	

## Appendix

1. Pregnancy-Related Deaths: Data from Maternal Mortality Review Committees in 36 US States, 2017–2019. Centers for Disease Control and Prevention. <https://www.cdc.gov/reproductivehealth/maternal-mortality/erase-mm/data-mmrc.html>. Accessed March 9, 2023.
2. Bauer ME, Lorenz RP, Bauer ST, Rao K, Anderson FWJ. Maternal deaths due to sepsis in the state of Michigan, 1999–2006. *Obstetrics and Gynecology*. 2015;126(4):747-752. doi:10.1097/AOG.0000000000001028
3. Kendle AM, Salemi JL, Tanner JP, Louis JM. Delivery-associated sepsis: trends in prevalence and mortality. *American Journal of Obstetrics and Gynecology*. 2019;220(4):391.e1-391.e16. doi:10.1016/j.ajog.2019.02.002
4. Sepsis in Obstetric Care. Alliance for Innovation on Maternal Health. <https://saferbirth.org/psbs/sepsis-in-obstetric-care/>. Accessed February 22, 2023.
5. Pareto Chart. Institute for Healthcare Improvement. <http://www.ihl.org:80/resources/Pages/Tools/ParetoDiagram.aspx>. Accessed April 13, 2022.
6. Priority Matrix: An Overlooked Gardening Tool. Institute for Healthcare Improvement. <http://www.ihl.org:80/education/IHIOpenSchool/resources/Pages/AudioandVideo/Priority-Matrix-An-Overlooked-Gardening-Tool.aspx>. Accessed April 13, 2022.
7. Impact Effort Matrix. American Society for Quality. <https://asq.org/quality-resources/impact-effort-matrix>. Accessed April 13, 2022.
8. Bauer ME, Housey M, Bauer ST, et al. Risk factors, etiologies, and screening tools for sepsis in pregnant women: a multicenter case-control study. *Anesthesia and Analgesia*. 2019;129(6):1613-1620. doi:10.1213/ANE.0000000000003709
9. Gibbs R, Bauer M, Olvera L, Sakowski C, Cape V, Main E. Improving diagnosis and treatment of maternal sepsis: a quality improvement toolkit. Stanford, CA: California Maternal Quality Care Collaborative.
10. Sepsis in Obstetrical Care: Element Implementation Details. Alliance for Innovation on Maternal Health. [https://saferbirth.org/wp-content/uploads/AIM\\_Bundle-EID\\_-SOC\\_Final\\_V1\\_2022.pdf](https://saferbirth.org/wp-content/uploads/AIM_Bundle-EID_-SOC_Final_V1_2022.pdf). Published online 2022.

11. Sepsis Bundle Project (SEP) National Hospital Inpatient Quality Measures. Centers for Medicare & Medicaid Services. [https://qualitynet.cms.gov/files/6391e95676962e0016ad9199?filename=2a-b\\_SEP-List\\_v5.14.pdf](https://qualitynet.cms.gov/files/6391e95676962e0016ad9199?filename=2a-b_SEP-List_v5.14.pdf). Accessed February 22, 2023.
12. Albright CM, Has P, Rouse DJ, Hughes BL. Internal validation of the sepsis in obstetrics score to identify risk of morbidity from sepsis in pregnancy. *Obstetrics and Gynecology*. 2017;130(4):747-755. doi:10.1097/AOG.0000000000002260
13. Urgent Maternal Warning Signs. Alliance for Innovation on Maternal Health. <https://saferbirth.org/aim-resources/aim-cornerstones/urgent-maternal-warning-signs/>. Accessed February 22, 2023.
14. How to Spot Sepsis. The UK Sepsis Trust. <https://sepsistrust.org/about/about-sepsis/>. Accessed February 22, 2023.
15. What is Sepsis? End Sepsis: The Legacy of Rory Staunton. <https://www.endsepsis.org/what-is-sepsis/>. Accessed February 22, 2023.
16. Pregnancy & Childbirth. Sepsis Alliance. <https://www.sepsis.org/sepsisand/pregnancy-childbirth/>. Accessed February 22, 2023.
17. Shields A, de Assis V, Halscott T. Top 10 pearls for the recognition, evaluation, and management of maternal sepsis. *Obstetrics and Gynecology*. 2021;138(2):289-304. doi:10.1097/AOG.0000000000004471
18. Maternal Safety Bundle for Sepsis in Pregnancy. Safe Motherhood Initiative. American College of Obstetricians and Gynecologists. <https://www.acog.org/-/media/project/acog/acogorg/files/forms/districts/smi-sepsis-bundle.pdf>. Accessed February 22, 2023.
19. UK Obstetric Surveillance System (UKOSS). NPEU: SHEER. <https://www.npeu.ox.ac.uk/ukoss>. Accessed February 22, 2023.
20. Three Ways to Create Psychological Safety in Health Care. Institute for Healthcare Improvement. <https://www.ihl.org:443/education/IHIOpenSchool/resources/Pages/AudioandVideo/Amy-Edmondson-Three-Ways-to-Create-Psychological-Safety-in-Health-Care.aspx>. Accessed February 22, 2023.
21. Frankel A, Haraden C, Lenoci-Edwards J. A Framework for Safe, Reliable, and Effective Care. Institute for Healthcare Improvement. [https://www.ihl.org/resources/Pages/IHIWhitePapers/Framework-Safe-Reliable-Effective-Care.aspx?PostAuthRed=/resources/\\_layouts/download.aspx?SourceURL=/resources/Knowledge%20Center%20Assets/IHIWhitePapers%20-%20AFrameworkforSafeReliableandEffectiveCare\\_d9247dfc-09ac-4777-bef1-62c7c9bd604c/FrameworkSafeReliableEffectiveCareWhitePaper.pdf](https://www.ihl.org/resources/Pages/IHIWhitePapers/Framework-Safe-Reliable-Effective-Care.aspx?PostAuthRed=/resources/_layouts/download.aspx?SourceURL=/resources/Knowledge%20Center%20Assets/IHIWhitePapers%20-%20AFrameworkforSafeReliableandEffectiveCare_d9247dfc-09ac-4777-bef1-62c7c9bd604c/FrameworkSafeReliableEffectiveCareWhitePaper.pdf). Accessed February 22, 2023.

22. Hear Her Campaign. Centers for Disease Control and Prevention. <https://www.cdc.gov/hearher/index.html>. Accessed February 22, 2023.
23. TeamSTEPS®. Agency for Healthcare Research and Quality. <https://www.ahrq.gov/teamsteps/index.html>. Accessed February 22, 2023.
24. Prevention of Infection After Gynecologic Procedures. American College of Obstetricians and Gynecologists. <https://www.acog.org/en/clinical/clinical-guidance/practice-bulletin/articles/2018/06/prevention-of-infection-after-gynecologic-procedures>. Accessed February 22, 2023.
25. Acosta CD, Kurinczuk JJ, Lucas DN, et al. Severe maternal sepsis in the UK, 2011–2012: a national case-control study. *PLOS Medicine*. 2014;11(7):e1001672. doi:10.1371/journal.pmed.1001672
26. Mhyre JM, D’Oria R, Hameed AB, et al. The maternal early warning criteria: a proposal from the national partnership for maternal safety. *Obstetrics and Gynecology*. 2014;124(4):782-786. doi:10.1097/AOG.0000000000000480
27. Shields LE, Wiesner S, Klein C, Pelletreau B, Hedriana HL. Use of maternal early warning trigger tool reduces maternal morbidity. *American Journal of Obstetrics and Gynecology*. 2016;214(4):527.e1-527.e6. doi:10.1016/j.ajog.2016.01.154
28. Committee Opinion No. 712 Summary: Intrapartum Management of Intraamniotic Infection. *Obstetrics and Gynecology*. 2017;130(2):490-492. doi:10.1097/AOG.0000000000002230
29. Stop. Look. Listen! Rutgers Robert Wood Johnson Medical School. [https://www.rwjms.rutgers.edu/RURWJ\\_SSLAnmatedPDF\\_FIN/document.pdf](https://www.rwjms.rutgers.edu/RURWJ_SSLAnmatedPDF_FIN/document.pdf). Accessed February 22, 2023.
30. Hear Her Campaign: Urgent Maternal Warning Signs. Centers for Disease Control and Prevention. <https://www.cdc.gov/hearher/maternal-warning-signs/index.html>. Accessed February 22, 2023.
31. SAVE YOUR LIFE: Get Care for These POST-BIRTH Warning Signs. Association of Women’s Health, Obstetric, and Neonatal Nurses. <https://www.awhonn.org/wp-content/uploads/2020/02/pbwssylhandoutenglish.pdf>. Accessed February 22, 2023.
32. POST-BIRTH Warning Signs Education Program. Association of Women’s Health, Obstetric, and Neonatal Nurses. <https://www.awhonn.org/education/hospital-products/post-birth-warning-signs-education-program/>. Accessed February 22, 2023.

33. Albright C, McCartney S, Hitti J. Identification and Management of Sepsis in Pregnancy: Obstetric Consensus Statement. [Identification-and-Management-of-Sepsis-in-Pregnancy-OB-Consensus-Statement.pdf \(uwmedicine.org\)](#). Accessed February 22, 2023.
34. Plante LA, Pacheco LD, Louis JM. SMFM consult series #47: sepsis during pregnancy and the puerperium. *American Journal of Obstetrics & Gynecology*. 2019;220(4):B2-B10. doi:10.1016/j.ajog.2019.01.216
35. Levels of maternal care: obstetric care consensus no, 9. *Obstetrics and Gynecology*. 2019;134(2):e41-e55. doi:10.1097/AOG.0000000000003383
36. ACOG Practice Bulletin No. 106: Intrapartum fetal heart rate monitoring: nomenclature, interpretation, and general management principles. *Obstetrics and Gynecology*. 2009;114(1):192-202. doi:10.1097/AOG.0b013e3181aef106
37. Post-Sepsis Syndrome. Sepsis Alliance. <https://www.sepsis.org/sepsis-basics/post-sepsis-syndrome/>. Accessed February 22, 2023.
38. Screening for Perinatal Depression. American College of Obstetricians and Gynecologists. <https://www.acog.org/en/clinical/clinical-guidance/committee-opinion/articles/2018/11/screening-for-perinatal-depression>. Accessed February 22, 2023.
39. Primary Care PTSD Screen for DSM-5 (PC-PTSD-5). Health Resources & Services Administration. <https://www.hrsa.gov/behavioral-health/primary-care-ptsd-screen-dsm-5-pc-ptsd-5>. Accessed February 22, 2023.
40. Health MC for WM. Screening for Perinatal Anxiety Using PASS. MGH Center for Women's Mental Health. <https://womensmentalhealth.org/posts/screening-for-perinatal-anxiety-using-pass-the-perinatal-anxiety-screening-scale/>. Accessed February 22, 2023.
41. National Maternal Mental Health Hotline. Maternal and Child Health Bureau. Health Resources & Services Administration. <https://mchb.hrsa.gov/national-maternal-mental-health-hotline>. Accessed February 22, 2023.
42. PostICU Resources. PostICU. <https://www.posticu.org/posticu-resources>. Accessed February 22, 2023.
43. Prescott HC, Angus DC. Enhancing recovery from sepsis: a review. *JAMA*. 2018;319(1):62-75. doi:10.1001/jama.2017.17687

44. Hardeman RR, Kheyfets A, Mantha AB, et al. Developing tools to report racism in maternal health for the cdc maternal mortality review information application (MMRIA): findings from the MMRIA racism & discrimination working group. *Maternal and Child Health Journal*. 2022;26(4):661-669. doi:10.1007/s10995-021-03284-3
45. Sun M, Oliwa T, Peek ME, Tung EL. Negative patient descriptors: documenting racial bias in the electronic health record. *Health Affairs*. 2022;41(2):203-211. doi:10.1377/hlthaff.2021.01423
46. Science of Improvement: Setting Aims. Institute for Healthcare Improvement.  
<https://www.ihl.org:443/resources/Pages/HowtoImprove/ScienceofImprovementSettingAims.aspx>. Accessed February 23, 2023.
47. Science of Improvement: Tips for Setting Aims. Institute for Healthcare Improvement.  
<https://www.ihl.org:443/resources/Pages/HowtoImprove/ScienceofImprovementTipsforSettingAims.aspx>. Accessed February 23, 2023.
48. Applying an Equity Lens to Performance Improvement. Institute for Healthcare Improvement.  
<https://www.ihl.org/communities/blogs/applying-an-equity-lens-to-performance-improvement>. Accessed February 23, 2023.