



ALLIANCE FOR INNOVATION
ON MATERNAL HEALTH



Perinatal Mental Health Conditions Patient Safety Bundle

Core Data Collection Plan
Version 1.1 January 2024



Perinatal Mental Health Conditions Patient Safety Bundlev

Core Data Collection Plan

Measurement Statement: For the purposes of this bundle, perinatal mental health conditions refer to mood, anxiety, and anxiety-related disorders that occur during pregnancy or within one year of delivery and are inclusive of mental health conditions with onset that predates pregnancy. These conditions include and are not limited to depression, anxiety and anxiety-related disorders like posttraumatic stress disorder and obsessive-compulsive disorder, bipolar disorder, and postpartum psychosis (see the [AIM PMHC ICD-10 Codes List](#) at the bottom of this document).

Although the PMHC patient safety bundle encompasses the entire perinatal period, the accompanying data collection plan primarily focuses on the inpatient birth admission due to the lack of universally accessible data pertaining to prenatal and postpartum care. In addition, these measures focus primarily on quality improvement opportunities within the influence of the inpatient care team. Suggested optional outpatient measures are included in the [Optional Measures](#) section.

State Surveillance Measures

Metric	Name	Description	Notes
PMHC SS1	Perinatal Mental Health Conditions (PMHC) among Pregnant and Postpartum People	<p>Report N/D <i>Disaggregate by race and ethnicity, payor</i></p> <p>Denominator: All qualifying pregnant and postpartum people during their birth admission</p> <p>Numerator: Among the denominator, those with any diagnosis of PMHC</p>	Use the severe maternal morbidity denominator criteria, subset by patients with a diagnosis of PMHC, to calculate the denominator for this measure.
PMHC SS2	Severe Maternal Morbidity among People with Perinatal Mental Health Conditions (excluding transfusion codes alone)	<p>Report N/D <i>Disaggregate by race and ethnicity, payor</i></p> <p>Denominator: All qualifying pregnant and postpartum people during their birth admission with PMHC</p> <p>Numerator: Among the denominator, those who experienced severe maternal morbidity, excluding those who experienced transfusion alone</p>	

Outcome Measures

Metric	Name	Description	Notes
PMHC O1	Percent of Pregnant and Postpartum People with PMHC Who Received or Were Referred to Treatment	<p>Report N/D Disaggregate by race and ethnicity, payor</p> <p>Denominator: Pregnant and postpartum people with a diagnosis of PMHC</p> <p>Numerator: Among the denominator, those with documentation of having received or been referred to treatment[§] prior to discharge from their birth hospitalization</p>	<p>[§]Treatment may include pharmacotherapy and/ or behavioral health therapy.</p> <p>Include in the numerator:</p> <ul style="list-style-type: none"> • Those who received treatment at any point during their pregnancy, regardless of current utilization • Those who did not receive treatment during pregnancy, but were referred to treatment prior to discharge from birth hospitalization

Process Measures

Metric	Name	Description	Notes
ALL P1- Version 2*	Provider and Nursing Education on Respectful and Equitable Care	<p>Report estimate in 10% increments (Round up)</p> <p>At the end of this reporting period, what cumulative proportion of OB clinicians[†] has received in the last 2 years an education on respectful and equitable care?</p>	<p>[†]The overarching intention of this measure is to capture all clinicians who work in a primarily inpatient OB service line or on an L&D, Antepartum, Postpartum unit. These clinicians will likely be interdisciplinary and could be inclusive of, but not limited to, nurses and nurse managers, advance practice nurses, nurse midwives, physician associates, and Family Medicine physicians or other specialties with delivering privileges at your institution.</p>

* This measure appears in other patient safety bundle data collection plans and are also referred to as multi-bundle measures. For the purposes of collecting data and reporting to the AIM Data Center, please collect and report this measure once per reporting period, regardless of the number of times they appear across data collection plans.

Metric	Name	Description	Notes
PMHC P1	Provider and Nursing Education on Perinatal Mental Health Conditions	<p>Report estimate in 10% increments (Round up)</p> <p>At the end of this reporting period, what cumulative proportion of OB clinicians[†] has received in the last 2 years education on care for pregnant and postpartum people with perinatal mental health conditions?</p>	<p>[†]The overarching intention of this measure is to capture all clinicians who work in a primarily inpatient OB service line or on an L&D, Antepartum, Postpartum unit. These clinicians will likely be interdisciplinary and could be inclusive of, but not limited to, nurses and nurse managers, advance practice nurses, nurse midwives, physician associates, and Family Medicine physicians or other specialties with delivering privileges at your institution.</p>
PMHC P2	Patient Education on Perinatal Mental Health Conditions	<p>Sample patient charts or report for all patients; report N/D <i>Disaggregate by race and ethnicity, payor</i></p> <p>Denominator: All pregnant and postpartum people during their birth admission, whether from sample or entire population</p> <p>Numerator: Among the denominator, those with documentation of having received verbal and written education on perinatal mental health conditions and when to seek care before discharge[‡]</p>	<p>For sampling guidance, please refer to the AIM Sampling Workbook.</p> <p>[‡]To be included in the numerator, patient record needs to include documentation of verbal and written education.</p>

Structure Measures

Metric	Name	Description	Notes
ALL S4*	Patient Education Materials on Urgent Postpartum Warning Signs	<p>Rate progress (1, not yet started – 5, fully in place) towards putting and keeping the structure measure fully in place</p> <p>Has your department developed/ curated patient education materials on urgent postpartum warning signs that align with culturally and linguistically appropriate standards?</p>	
ALL S6*	Inpatient-Outpatient Care Coordination Workgroup	<p>Rate progress (1, not yet started – 5, fully in place) towards putting and keeping the structure measure fully in place</p> <p>Has your hospital established a multidisciplinary workgroup of inpatient and outpatient providers that meets regularly to identify and implement best practices on issues related to pregnancy and the postpartum period that cross the continuum of care?</p>	<ul style="list-style-type: none"> • This workgroup should help coordinate the completion of the other structure measures
ALL S7*	Resource Mapping/ Identification of Community Resources	<p>Rate progress (1, not yet started – 5, fully in place) towards putting and keeping the structure measure fully in place</p> <p>Has your hospital created a comprehensive list of community resources, customized to include resources relevant for pregnant and postpartum people, that will be shared with all postpartum inpatient nursing units and outpatient OB sites?</p>	<ul style="list-style-type: none"> • Resource list should be updated annually • Resource list should include OUD/SUD treatment resources as well as mental health resources and allow for customization based on patient population (e.g., BIPOC).

* This measure appears in other patient safety bundle data collection plans and are also referred to as multi-bundle measures. For the purposes of collecting data and reporting to the AIM Data Center, please collect and report this measure once per reporting period, regardless of the number of times they appear across data collection plans.

Metric	Name	Description	Notes
PMHC S1	Perinatal Mental Health Assessment and Response Protocol	<p>Rate progress (1, not yet started – 5, fully in place) towards putting and keeping the structure measure fully in place</p> <p>Does you have a written assessment and response protocol for perinatal mental health conditions that is tiered based on illness severity and risk of harm?</p>	
PMHC S2	Validated PMHC Screening Tools Shared with Prenatal Care Sites	<p>Rate progress (1, not yet started – 5, fully in place) towards putting and keeping the structure measure fully in place</p> <p>Has your hospital shared with all its affiliated prenatal care sites validated screening tools for diagnosis of PMHC?</p>	

Optional Measures

Metric	Name	Description	Notes
PMHC OP1	Proportion of Pregnancy-Associated Deaths Due to Suicide	<p>Report N/D <i>Disaggregate by race and ethnicity, payor</i></p> <p>Denominator: Total pregnancy-associated deaths</p> <p>Numerator: Among the denominator, pregnancy-associated deaths due to suicide</p>	

Metric	Name	Description	Notes
PMHC OP2	Proportion of Pregnancy-Associated Deaths Due to Overdose	<p>Report N/D <i>Disaggregate by race and ethnicity, payor</i></p> <p>Denominator: Total pregnancy-associated deaths</p> <p>Numerator: Among the denominator, pregnancy-associated deaths due to overdose</p>	This optional measure is included in acknowledgement that intention of overdose is not always known or able to be determined by maternal mortality review committees
PMHC OP3	Prenatal PMHC Screening	<p>Sample patient charts or report for all patients; report N/D for each sub measure <i>Disaggregate by race and ethnicity, payor</i></p> <p>OP3A. Denominator: All patients with at least one prenatal visit during the reporting period</p> <p>Numerator: Among the denominator, patients who were screened at least once for depression using a validated tool</p> <p>OP3B. Denominator: All patients with at least one prenatal visit during the reporting period</p> <p>Numerator: Among the denominator, patients who were screened at least once for anxiety using a validated tool</p>	<ul style="list-style-type: none"> • For sampling guidance, please refer to the AIM Sampling Workbook • If using a single tool valid for both depression and anxiety, report the same statistic for both OP3A and OP3B • These measures are not intended to be used in place of related HEDIS® measures for prenatal depression screening and follow up. These measures may be used by outpatient teams to drive quality improvement efforts with the goal of improving patient care and HEDIS® measure outcomes

Metric	Name	Description	Notes
PMHC OP4	Postpartum PMHC Screening	<p>Sample patient charts or report for all patients; report N/D <i>Disaggregate by race and ethnicity, payor</i></p> <p>OP4A. Denominator: All patients with at least one postpartum visit during the reporting period</p> <p>Numerator: Among the denominator, patients who were screened at least once for depression using a validated tool</p> <p>OP4B. Denominator: All patients with at least one postpartum visit during the reporting period</p> <p>Numerator: Among the denominator, patients who were screened at least once for anxiety using a validated tool</p>	<ul style="list-style-type: none"> • For sampling guidance, please refer to the AIM Sampling Workbook • If using a single tool valid for both depression and anxiety, report the same statistic for both OP4A and OP4B • These measures are not intended to be used in place of related HEDIS® measures for postpartum depression screening and follow up. These measures may be used by outpatient teams to drive quality improvement efforts with the goal of improving patient care and HEDIS® measure outcomes

AIM PMHC ICD-10 Codes List

Condition Group	Definition
Depression	F320, F321, F322, F323, F324, F325, F328, F3289, F329, F32A, F330, F331, F332, F333, F334, F3340, F3341, F3342, F338, F339, F341, F530, O906, O9934
Bipolar disorder	F3010, F3011, F3012, F3013, F302, F303, F304, F308, F309, F310, F3110, F3111, F3112, F3113, F312, F3130, F3131, F3132, F314, F315, F3160, F3161, F3162, F3163, F3164, F3170, F3171, F3172, F3173, F3174, F3175, F3176, F3177, F3178, F3181, F3189, F319, F340, F3481, F3489, F39
Anxiety	F064, F409, F4000, F4001, F4002, F4010, F4011, F40218, F40240, F40241, F408, F410, F411, F413, F418, F419, F430, F458, F488, F489, F938, F99, R457
Posttraumatic stress disorder	F4310, F4311, F4312
Obsessive-compulsive disorder	F422, F423, F424, F428, F429, R4681
Psychosis	F060, F062, F200, F201, F202, F203, F205, F2081, F2089, F209, F21, F22, F23, F24, F250, F251, F258, F259, F28, F29, F531, F440, F441, F442, F4481, F4489, F449, F481, F482
Other	R45850, R45851, F061, F0630, F0631, F0632, F0633, F0634, F349, F4320, F4321, F4322, F4323, F4324, F4325, F4329, F4381, F4389, F439, F444, F445, F446, F447, F450, F451, F4520, F4521, F4522, F4529, F4541, F4542, F458, F459, F5101, F5102, F5103, F5104, F5105, F5109, F5111, F5112, F5113, F5119, F518, F519, F54, F59, O99340, O99341, O99342, O99343, O99344, O99345

This document was developed with support by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) as part of an award, UC4MC49476, totaling \$3,000,000 with 0% financed with non-governmental sources. The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement, by HRSA, HHS, or the U.S. Government. For more information, please visit [HRSA.gov](https://www.hrsa.gov).

© 2023 American College of Obstetricians and Gynecologists. Permission is hereby granted for duplication and distribution of this document, in its entirety and without modification, for solely non-commercial activities that are for educational, quality improvement, and patient safety purposes. All other uses require written permission from ACOG.